

# Altermagnetism: The Case of MnTe

R. D. Gonzalez Betancourt<sup>1</sup>, J. Krempaský<sup>2</sup>, S. W. D'Souza<sup>3</sup>, J. Zubáč<sup>1</sup>,  
R. J. Gonzalez-Hernandez<sup>4</sup>, A. Hariki<sup>5</sup>, Z. Šobán<sup>1</sup>, G. Springholz<sup>6</sup>, K. Uhlířová<sup>7</sup>,  
K. Olejník<sup>1</sup>, J. Kuneš<sup>8</sup>, J. Minár<sup>3</sup>, J. H. Dil<sup>9</sup>, K. Výborný<sup>1</sup>, J. Sinova<sup>10</sup>,  
T. Jungwirth<sup>1,11</sup>, P. Wadely<sup>11</sup>, S. T. B. Goennenwein<sup>12</sup>, A. Thomas<sup>13</sup>,  
H. Reichlová<sup>1</sup>, J. Železný<sup>1</sup>, L. Šmejkal<sup>1,10</sup>, D. Kriegner<sup>1, a)</sup>

<sup>1</sup>*Institute of Physics ASCR, Praha, Czech Republic*

<sup>2</sup>*Photon Science Division, Paul Scherrer Institut, Villigen, Switzerland*

<sup>3</sup>*University of West Bohemia, New Technologies Research Center, Plzen, Czech Republic*

<sup>4</sup>*Grupo de Investigación en Física Aplicada, Departamento de Física, Universidad del Norte, Barranquilla, Colombia*

<sup>5</sup>*Department of Physics and Electronics, Graduate School of Engineering, Osaka Metropolitan University, Osaka, Japan*

<sup>6</sup>*Institute of Semiconductor and Solid State Physics, Johannes Kepler University Linz, Austria*

<sup>7</sup>*Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic*

<sup>8</sup>*Institute for Solid State Physics, TU Wien, 1040 Vienna, Austria*

<sup>9</sup>*Institut de Physique, École Polytechnique Fédérale de Lausanne, Switzerland*

<sup>10</sup>*Institut für Physik, Johannes Gutenberg Universität Mainz, Germany*

<sup>11</sup>*School of Physics and Astronomy, University of Nottingham, United Kingdom*

<sup>12</sup>*Department of Physics, University of Konstanz, Germany*

<sup>13</sup>*Leibniz Institute of Solid State and Materials Research (IFW), Dresden, Germany*

<sup>a)</sup> Corresponding author: kriegner@fzu.cz

**Abstract.** Altermagnets emerge as a novel type of compensated collinear magnets that complement the conventional classes of ferro- and antiferromagnets [1,2]. In contrast to antiferromagnets, for which the opposite sublattices are connected by simple lattice translation or inversion, in altermagnets, they are connected by a lattice rotational symmetry (symmorphic/non-symmorphic, proper/improper) [1,2]. This causes time reversal symmetry breaking in the band structure and – in contrast to ferromagnets – a spin splitting that alternates in sign within the Brillouin zone suggesting the term altermagnetism. The magnetization, however, integrates to zero over the entire Brillouin zone. Here we show experimental verification of the spin-splitting by angle resolved photo emission spectroscopy investigations in MnTe [3]. The characteristic spin splitting further enables linear responses such as the anomalous Hall effect [4-6], and X-ray magnetic circular dichroism [7] which will be also discussed.

## REFERENCES

- [1] L. Šmejkal, J. Sinova, and T. Jungwirth, *Phys. Rev. X* **12**, 031042 (2022)
- [2] L. Šmejkal, J. Sinova, and T. Jungwirth, *Phys. Rev. X* **12**, 040501 (2022)
- [3] J. Krempaský, *et al.*, *Nature* **626**, 517-522 (2024)
- [4] R. D. Gonzalez Betancourt, *et al.*, *Phys. Rev. Lett.* **130**, 036702 (2023)
- [5] T. Tschirner, *et al.*, *APL Mater.* **11**, 101103 (2023)
- [6] H. Reichlova, *et al.*, *Nat. Comm.* (accepted) (2024)
- [7] A. Hariki, *et al.*, *Phys. Rev. Lett.* (accepted) (2024)